

BOOK OF PROCEEDINGS

**Fifth International Scientific Agricultural Symposium
“Agrosym 2014”**

AGROSYM 2014



Jahorina, October 23 - 26, 2014

Impressum

Fifth International Scientific Agricultural Symposium „Agrosym 2014“

Book of Proceedings

Published by

University of East Sarajevo, Faculty of Agriculture, Republic of Srpska, Bosnia
University of Belgrade, Faculty of Agriculture, Serbia
Mediterranean Agronomic Institute of Bari (CIHEAM - IAMB) Italy
International Society of Environment and Rural Development, Japan
Balkan Environmental Association, B.EN.A, Greece
Academy of Engineering Sciences of Serbia, Serbia
Maize Research Institute „Zemun Polje“ Serbia
Biotechnical Faculty, University of Montenegro, Montenegro
Balkan Scientific Association of Agricultural Economics, Serbia
Institute of Agricultural Economics, Serbia
Faculty of Agriculture, University of Banja Luka, Bosnia and Herzegovina

Editor in Chief

Dusan Kovacevic

Technical editors

Sinisa Berjan
Milan Jugovic
Mirjana Radovic
Noureddin Driouech
Rosanna Quagliariello

Website:

<http://www.agrosym.rs.ba>

CIP - Каталогизacija у публикацији
Народна и универзитетска библиотека
Републике Српске, Бања Лука

631(082)(0.034.2)

INTERNATIONAL Scientific Agricultural Symposium "Agrosym
2014" (5 ; Jahorina)

Book of proceedings [Elektronski izvor] / Fifth International
Scientific Agricultural Symposium "Agrosym 2014", Jahorina,
October 23 - 26, 2014 ; [editor in chief Dušan Kovačević]. - East
Sarajevo = Istočno Sarajevo : Faculty of Agriculture = Poljoprivredni
fakultet, 2014. - 1 elektronski optički disk (CD-ROM) : tekst, slika ;
12 cm

CD ROM čitač. - Dostupno i na:
http://www.agrosym.rs.ba/agrosym/agrosym_2014/index.html. -
Nasl. sa nasl. ekrana. - Bibliografija uz svaki rad. - Registar.

ISBN 978-99955-751-9-9

COBISS.RS-ID 4641816

**Fifth International Scientific Agricultural Symposium “Agrosym 2014”
Jahorina, October 23-26, 2014, Bosnia and Herzegovina**

HONORARY COMMITTEE

STEVO MIRJANIC, Minister of Agriculture, Water Management and Forestry of Republic of Srpska, Bosnia; JASMIN KOMIC, Minister of Science and Technology of Republic of Srpska, Bosnia; GORAN MUTABDZIJA, Minister of Education and Culture of Republic of Srpska, Bosnia; RADOSLAV GRUJIC, Rector of the University of East Sarajevo, Bosnia; MILICA PETROVIC, Dean of the Faculty of Agriculture, University of Belgrade, Serbia; COSIMO LACIRIGNOLA, Director of the Mediterranean Agronomic Institute of Bari, Italy; MARIO T. TABUCANON, President of the International Society of Environment and Rural Development, Japan; FOKIAON K. VOSNIAKOS, President of the Balkan Environmental Association (B.EN.A), Greece; BRANKO KOVACEVIC, President of the Academy of Engineering Sciences of Serbia, Serbia; BRANKA KRESOVIC, Director of the Maize Research Institute “Zemun Polje”, Serbia; MIOMIR JOVANOVIC, Dean of the Biotechnical Faculty, University of Podgorica; DRAGO CVIJANOVIC, Director of the Institute of Agricultural Economics, Serbia; NIKOLA MICIC, Dean of the Faculty of Agriculture, University of Banja Luka, Bosnia.

SCIENTIFIC COMMITTEE

DUSAN KOVACEVIC, Faculty of Agriculture, University of Belgrade, Serbia; WILLIAM MEYERS, Howard Cowden Professor of Agricultural and Applied Economics, University of Missouri, USA; JOHN BRAYDEN, Norwegian Agricultural Economics Research Institute (NILF), Norway; STEVE QUARIE, Visiting Professor, School of Biology, Newcastle University, United Kingdom; ATEF HAMDY, Emeritus Professor, Land and Water Resources Department; IAMB, Italy; DANI SHTIENBERG, full professor, Department of Plant pathology and Weed Research, ARO, the Volcani Center, Bet Dagan, Israel; THOMAS G. JOHNSON, University of Missouri – Columbia, USA; DIETER TRAUTZ, University of Applied Science, Germany; MACHITO MIHARA, Tokyo University of Agriculture, Japan; MARKUS SCHERMER, Department of Sociology, University of Innsbruck, Austria; FOKIAON VOSNIAKOS, Balkan Environmental Association (B.EN.A), Greece; MATTEO VITTUARI, Faculty of Agriculture, University of Bologna, Italy; VELIBOR SPALEVIC, The journal “Agriculture and Forestry”, Biotechnical Faculty, University of Montenegro, Montenegro; EPN UDAYAKUMARA, Faculty of Applied Sciences, Sabaragamuwa University, Sri Lanka; VLADIMIR SMUTNÝ, full professor, Mendel University, Faculty of agronomy, Czech Republic; FRANC BAVEC, full professor, Faculty of Agriculture and Life Sciences, Maribor, Slovenia; NICOLAE ESTUDOE, full professor, Academy of Economic Studies, Bucharest, Romania; JAN MOUDRÝ, full professor, Faculty of Agriculture, South Bohemia University, Czech Republic; STEFAN TYR, full professor, Faculty of Agrobiology and Food Resources, Slovakia; NATALIJA BOGDANOV, Faculty of Agriculture, University of Belgrade, Serbia; SABAHUDIN BAJRAMOVIC, Faculty of Agriculture and Food Sciences, University of Sarajevo, Bosnia; FRANCESCO PORCELLI, University of Bari Aldo Moro, Italy; VASILJE ISAJEV, Faculty of Forestry, University of Belgrade, Serbia; ELAZAR FALLIK, Agricultural Research Organization (ARO), Volcani, Israel; JUNAID ALAM MEMON, Pakistan Institute of Development Economics, Pakistan; HIROMU OKAZAWA, Faculty of Regional Environment Science, Tokyo University of Agriculture, Japan; MLADEN TODOROVIC, Land and Water Resources Department; IAMB, Italy; HAMID EL BILALI, Sustainable Agriculture, Food and Rural Development Department. IAMB, Italy; NOUREDDIN DRIUECH, Environmental Sciences and Organic Agriculture, IAMB, Italy; LALITA SIRIWATTANANON, Faculty of Agricultural Technology, Rajamangala University of Technology Thanyaburi (RMUTT), Thailand; ABID HUSSAIN, International Centre for Integrated Mountain Development (ICIMOD), Nepal; AMRITA GHATAK, Gujarat Institute of Development Research (GIDR), India; MÁRTA BIRKÁS, full professor, St. Istvan University, Godollo – Hungary; UDAI PRATAP SINGH, Department of Mycology and Plant Pathology, Banaras Hindu University, India; ANDRZEJ KOWALSKI, Director of the Institute for Agricultural and Food Economy, Warszawa-Poland; YALCIN KAYA, The Director of the Plant Breeding Research Center, University of Trakya, Turkey; KOSANA KONSTATINOV, Academy of Engineering Sciences of Serbia, Serbia; NEBOJSA MOMIROVIC, Faculty of Agriculture, University of Belgrade, Serbia; ZORAN JOVOVIC, Biotechnical Faculty, University of Montenegro, Montenegro; VLADIMIR VUKADINOVIC, full professor, Faculty of Agriculture, University of Osijek, Croatia; DANIJEL JUG, associate professor, Faculty of Agriculture, University of Osijek, Croatia; VLADO KOVACEVIC, full professor, Faculty of Agriculture, University of Osijek, Croatia; MILAN MARKOVIC, Department for Animal husbandry, Biotechnical Faculty, University of Montenegro, Montenegro.

ORGANIZATION COMMITTEE

VESNA MILIC, Dean of the Faculty of Agriculture, University of East Sarajevo, Bosnia; DEJAN BOKONJIC, Vice rector of the University of East Sarajevo, Bosnia; STEVAN TRBOJEVIC, Vice rector of the University of East Sarajevo, Bosnia; PREDRAG PUDJA, Vice dean of the Faculty of Agriculture, University of Belgrade, Serbia; JEGOR MILADINOVIC, Scientific Advisor, Institute of Field and Vegetable Crops, Novi Sad, Serbia; ZELJKO DOLJANOVIC, Faculty of Agriculture, University of Belgrade, Serbia; ROBERTO CAPONE, Mediterranean Agronomic Institute of Bari, Italy; ROSANNA QUAGLIARIELLO, Mediterranean Agronomic Institute of Bari, Italy; NOUREDDIN DRIQUECH, Coordinator of MAIB Alumni Network (FTN), Mediterranean Agronomic Institute of Bari, Italy; ALEKSANDRA DESPOTOVIC, Biotechnical Faculty Podgorica, University of Montenegro, Montenegro; MILIC CUROVIC, The journal “Agriculture and Forestry”, Biotechnical Faculty Podgorica, University of Montenegro, Montenegro; GORAN PERKOVIC, Faculty of Agriculture, University of East Sarajevo, Bosnia; MIRJANA RADOVIC, Faculty of Agriculture, University of East Sarajevo, Bosnia; MILAN JUGOVIC, Faculty of Agriculture, University of East Sarajevo, Bosnia; SINISA BERJAN, Faculty of Agriculture, University of East Sarajevo, Bosnia, secretary

GRAIN YIELD, YIELD COMPONENTS AND MALTING QUALITY TRAITS OF SPRING BARLEY ON ACIDIC SOILS

Milomirka MADIC^{1*}, Dragan DJUROVIC¹, Aleksandar PAUNOVIC¹, Miodrag JELIC²,
Desimir KNEZEVIC², Branka GOVEDARICA³

¹University of Kragujevac, Faculty of Agronomy, Zemun, Serbia

²Faculty of Agriculture, Lešak, University of Priština, Kosovska Mitrovica

³University of East Sarajevo, Faculty of Agriculture

*Corresponding author: mmadic@kg.ac.rs

Abstract

A varietal trial involving six genotypes of spring barley ('Jastrebac', 'Dinarac', 'Dunavac', 'NS-404', 'Jaran' and 'Lider') was conducted over a three-year period at the farm estate of the Secondary School of Agriculture and Chemistry, Kraljevo (Serbia) (the experimental field of the Faculty of Agronomy, Zemun) on an acidic soil (pH_{H2O} 4.5). The following traits were analysed: stem height, spike length, grain number per spike, thousand-kernel weight, grain yield, total germination capacity, grain protein content and extract content. Regardless of year, stem height was significantly higher in 'Dunavac' and 'Jaran' than in the other cultivars. The highest stem height in all cultivars was obtained in the second year. Spike length, grain number per spike and 1000-kernel weight were significantly higher in 'NS-404' compared to the other cultivars. Variations in grain yield across years were the lowest in 'Jaran' and 'Dinarac'. Total germination capacity was significantly higher in 'Jastrebac' than in the other cultivars in the first and second years. The significant increase in protein content in both years on average for all cultivars was accompanied by a significant decrease in malt extract, and vice versa. The high protein content in some cultivars along with the high malt extract content suggests that the strength of correlation between the traits is dependent on environmental conditions. Although the soil had poor physical and chemical properties, all cultivars had their 1000-kernel weight, total germination capacity, protein content and malt extract content within the standardised limits.

Key words: *barley, grain yield, yield components, malting quality traits*

Introduction

The proper choice of cultivar is of special importance in obtaining high grain yields in good-quality malt barley. In addition, different soil and climate conditions, notably temperature and moisture content during grain fill (Passarella et al., 2005) can largely contribute to variations in major yield components and, hence, total yield and grain quality of malting barley (Atlin et al., 2000; Paunović et al., 2007; Madić et al., 2009). Pržulj et al. (2014) stressed that temperature and rainfall do not play the leading role in determining grain yield and quality of malting barley since grain quality is often poor when these factors approach optimal values. Moreover, grain yield and quality attributes in barley are affected by certain cultural practices such as nitrogen fertilisation (Pržulj and Momčilović, 2008; Marconi et al., 2010). As stressed by Pržulj and Momčilović (2002), realising the full yield and quality potential in barley necessitates strict adherence to production technology, given the precise quality requirements of the brewing industry for barley grain and malt. To ensure high quality of raw materials, new cultivars should be analysed for their traits and introduced into the production system, and cultivar choice made for a specific region. Grain quality i.e. malt extract content is largely dependent on growing conditions (temperature, available moisture, available N, fertilisation)