

RESEARCH ARTICLE

Mortality from Stomach Cancer in Serbia, Excluding the Province of Kosovo, in the 1991-2009 Period

Milena Ilic^{1*}, Tanja Prodic¹, Zoran Milosavljevic², Biljana Ljubic³

Abstract

Background/Aim: Stomach cancer is the second most common cause of death from all malignant tumors in the world (third in men, fifth in women), with a strong decreasing trend in most developed countries. The aim of this descriptive epidemiological study was to analyze mortality of stomach cancer in Serbia, excluding the Province of Kosovo, in the 1991-2009 period. **Materials and Methods:** In data analysis, we used mortality rates which were standardized directly using those of the world population as a standard. In order to analyze the mortality trend from stomach cancer, linear trend and regression analysis were used. Confidence intervals (CIs) for the average age-adjusted and age-specific mortality rates were assessed with 95% level of probability. Mortality data were derived from the data file of the Statistical Office of the Republic of Serbia. **Results:** During the 1991-2009 period, a significant downward trend in mortality of stomach cancer was recorded in Serbia ($y=9.78 - 0.13x$, $p=0.000$; average annual percent change was -6.3 (95% CI, -7.8 to -4.8). During the same period, a significant decrease in mortality trend was found both in male ($y=14.13 - 0.20x$; $p=0.000$; % change was -7.7 (95% CI, -10.9 to -4.5) and female populations ($y=6.27 - 0.08x$; $p=0.000$; % change was -4.4 (95% CI, -5.3 to -3.6). **Conclusion:** Decreasing trends in mortality from stomach cancer in Serbia are similar to those in most developed countries.

Keywords: Stomach cancer - mortality - descriptive epidemiologic study

Asian Pacific J Cancer Prev, 14 (3), 2067-2070

Introduction

Malignant stomach tumors are the second leading cause of death among malignant tumors, with more than 730,000 deaths, accounting for 10% of total deaths worldwide in 2008 (Jemal et al., 2011). The highest stomach cancer mortality rates in 2008 were recorded in Asia (Mongolia - 26.5 per 100,000 inhabitants, China - 22.3 per 100,000 inhabitants, Bhutan - 22.1 per 100,000 inhabitants) (IARC, 2008). The lowest mortality rates were recorded in certain South African countries (Botswana - 0.3 per 100,000 inhabitants and Namibia - 1.1 per 100,000 inhabitants), and United States of America - 2.0 per 100,000 inhabitants. Over 75% of stomach cancer cases (556,471 cases) occur in developing countries (353,685 men vs. 202,786 women) (IARC, 2008; Jemal et al., 2011). Nearly 72% of deaths were recorded in Asia, almost 66.5% of which occurred in China.

Stomach cancer rates are about twice as high in males as in females (Parkin, 2004; Ferlay et al., 2010; Jemal et al., 2011). Stomach cancer mortality starts to rise in middle-aged adults of both genders (45-64 years of age), and becomes even more frequent in the elderly (75 years of age and above) (Look et al., 2001). The mortality


of stomach cancer has decreased considerably in most developed countries over the past several decades (Levi et al., 2003). Similar trends, but with later onset, were also observed in some Asian countries, including Japan and China (Lee et al., 2006).

The aim of this descriptive epidemiological analysis was to estimate death rates for stomach cancer and their secular trends in the population of Serbia over the 1991-2009 period.

Materials and Methods

The mortality rates are calculated from data collected by the Statistical Office of the Republic of Serbia, which receives death certificates and compiles mortality data by gender, age, year, and cause of death. We used the International Classification of Diseases, 9th and 10th Revision, to classify codes of deaths (WHO). Data on the deceased from all causes of death were used in the study (codes 001-999 by 9th and A00-Z99 by 10th revision of the International Statistical Classification of Diseases, Injuries and Causes of Death), all malignant tumors (codes 140-209, 9th Revision and C00-C97, 10th Revision), symptoms, signs and abnormal clinical and laboratory findings (codes

¹Department of Epidemiology, ²Department of Histology, ³Center for Molecular Medicine and Stem Cell Research, Faculty of Medical Sciences, University of Kragujevac, Kragujevac, Serbia *For correspondence: drmilena@iic@yahoo.com



observation period, a significant decrease in the rates of deaths caused by stomach cancer is recorded in men aged 20-29, 30-39, 40-49, 50-59, 60-69, 70-79. The female population also demonstrated a significant decrease in deaths caused by stomach cancer in age groups with middle-aged (30-39 years of age) and elderly women (50-59, 60-69, 70-79). None of the regression analysis models corresponded to mortality data for other age groups.

Discussion

In spite of the strong downward mortality rates worldwide, stomach cancer is the second most important cause of death from cancer worldwide (Parkin, 2004; Jemal et al., 2011). In Serbia, with 1313 deaths and 5.4% in total number of deaths from malignant tumors in 2008, stomach cancer takes the fourth place.

Stomach cancer mortality rates in different populations around the world vary by more than tenfold (IARC, 2008). Nearly three-fourths of stomach cancers occur in developing countries. Mortality rates in men and women in 2008 ranged from 20.3 per 100,000 people in East Asia to 2.1 per 100,000 in North America (IARC, 2008; Ferlay et al., 2010). The highest mortality rates were recorded in Asia (Mongolia -26.5 per 100,000) and parts of South America (Honduras -22.3 per 100,000, Peru 18.2, Chile -15.0) in men and women. In contrast, the lowest rates were observed in men and women in North America and most parts of Africa (approximately 2 per 100,000). Across Europe, major differences in stomach cancer death rates are evident (IARC, 2008). The stomach cancer rates in Russian Federation (15.2 per 100,000) are more than five times higher than those in Scandinavia or France (close to 3.0 per 100,000), and higher stomach cancer rates are generally observed in Eastern Europe (Kyrgyzstan -20.1, Belarus -18.8). Moderately high death rates of stomach cancer are observed in Bosnia and Herzegovina (7.7 per 100,000), Serbia (7.4), Italy (6.6), Spain (6.0), Greece (5.0), Finland (4.4). The lowest rates in Europe in 2008 are found in Cyprus (2.9 per 100,000), Switzerland (3.0), and Iceland, Sweden and France metropolitan (3.2 per 100,000 all).

Mortality rates in men are 2-3 times higher than in women. The highest mortality rates are estimated in Eastern Asia (28.1 per 100,000 in men, 13.0 per 100,000 in women), and the lowest in Northern America (2.8 and 1.5, respectively). High mortality rates are also present in both genders in Central and Eastern Europe, and in Central and South America (IARC, 2008; Jemal et al., 2011). Differences in stomach cancer mortality in men and women are associated with differences in incidence of various adenocarcinoma subtypes by histology (intestinal/diffuse) and location (proximal/distal) (Munoz et al., 1968; Ekström et al., 2000). Diffuse adenocarcinoma occur more frequently in women, while in men is dominant intestinal adenocarcinoma, the subtype being responsible for most of the international variations.

Stomach cancer is rare in children and young adults. In Serbia, as in most countries, stomach cancer mortality rates start to rise in middle-aged adults (45-64 years of age), and it is even more common in the elderly (75

years of age and older) age group for males and females, respectively (Howlader et al., 2011). In the second half of 20th century, stomach cancer mortality in Japan has reached the mortality levels in Western countries, which may be explained by introduction of stomach cancer screening in Japan and life style changes, including reduced salt intake and increased intake of fruit and vegetables, reduction in smoking and prevention of *Helicobacter pylori* infection (Inoue et al., 2005; Lee et al., 2006). In addition, studies of immigrants suggest significant reduction of stomach cancer risk in Japanese immigrants to the lower risk areas (Maskarinec et al., 2004), even though numerous results indicate the importance of exposure to environmental factors during childhood in determining risk (Coggon et al., 1990).

In both men in women, in the past decades the stomach cancer mortality has been showing significant downward trend in most of the developed countries (Amiri et al., 2011; Howlader et al., 2011). Stomach cancer mortality rates declined between 1980 and 2005 at about the same rate (3.6-4.9% per year) for both men and women in seven European countries (Denmark, Finland, France metropolitan, the Netherlands, Norway, Sweden, and England and Wales) (Amiri et al., 2011). The US mortality trend for stomach cancer between 1990-2008, showed significant decrease for all races, with the average annual percent change -3.0% in both genders, and -3.4% in male and -2.7% in female (Howlader et al., 2011). Similar trend, but with later onset, was observed in Asian countries, including Japan and China (Lee et al., 2006; Yang, 2006).

The international variation in mortality, time trends, and the impact of migration on mortality from stomach cancer are mainly due to differences in socio-economic environment and lifestyle, education and health care system has begun in recent decades. Factors leading to the decrease in stomach cancer mortality include broader availability of fresh fruit and vegetables, less salt and conserved food in the diet, better food conservation including refrigeration, decreased incidence of chronic infections with *Helicobacter pylori* resulting from improved hygiene and use of antibiotics, and introduction of the screening program (Levi et al., 2004; Parkin, 2004; Lee et al., 2006; Yang, 2006). Evidence linking *Helicobacter pylori* infection with stomach cancer risk contributed to the 1994 International Agency for Research on Cancer's classification of the bacterium as carcinogenic to humans (IARC, 1994). International prevalence variations of *Helicobacter pylori* infection demonstrate certain similarities with those of the stomach cancer; in developing countries, prevalence of *Helicobacter pylori* infection in adults is estimated to 76% vs. 58% in developed countries. High prevalence of the *Helicobacter pylori* infection with minor international differences indicates that some other factors play significant role in the stomach cancer etiology (Parkin et al., 2002). While some authors (Ngoan et al., 2002; Ito et al., 2003) believe diet plays no role in the stomach cancer etiology, American Cancer Society lists smoked foods, salted fish and meats, and pickled vegetables as risk factors for stomach cancer (American Cancer Society, 2010). On the other hand, eating fresh fruits and vegetables that contain antioxidative

References

- American Cancer Society. Stomach cancer. 2010. Atlanta, Ga: American Cancer Society; 2010. Available at: <http://www.cancer.org/acs/groups/cid/documents/>
- Amiri M, Janssen F, Kunst AE (2011). The decline in stomach cancer mortality: exploration of future trends in seven European countries. *Eur J Epidemiol*, **26**, 23-8.
- Coggon D, Osmond C, Barker DJ (1990). Stomach cancer and migration within England and Wales. *Br J Cancer*, **61**, 573-4.
- Ekström AM, Hansson LE, Signorello LB, et al (2000). Decreasing incidence of both major histologic subtypes of gastric adenocarcinoma--a population-based study in Sweden. *Br J Cancer*, **83**, 391-6.
- Ferlay J, Shin HR, Bray F, et al (2010). Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *Int J Cancer*, **127**, 2893-917.
- Gołkowski F, Szybiński Z, Rachtan J, et al (2007). Iodine prophylaxis--the protective factor against stomach cancer in iodine deficient areas. *Eur J Nutr*, **46**, 251-6.
- Howlander N, Noone AM, Krapcho M, et al (eds). SEER Cancer Statistics Review, 1975-2008, National Cancer Institute. Bethesda, MD, http://seer.cancer.gov/csr/1975_2008/, based on November 2010 SEER data submission, posted to the SEER web site, 2011.
- IARC Monograph on the Evaluation of Carcinogenic Risks to Humans, Vol. 61. Schistosomes, Liver Flukes and *Helicobacter pylori*. Lyon, France: International Agency for Research on Cancer; 1994. Available at: <http://monographs.iarc.fr/>
- IARC Monographs on the Evaluation of Carcinogenic Risk to Humans. Vol. 83: Tobacco Smoke and Involuntary Smoking. Lyon, France: International Agency for Research on Cancer; 2004. Available at: <http://monographs.iarc.fr/>
- IARC. World Cancer Report, 2008. Boyle P. and Levin BE. (eds), IARC Press, Lyon. Available at: <http://monographs.iarc.fr/>
- Inoue M, S Tsugane S (2005). Epidemiology of gastric cancer in Japan. *Postgrad Med J*, **81**, 419-24.
- Ito LS, Inoue M, Tajima K, et al (2003). Dietary factors and the risk of gastric cancer among Japanese women: a comparison between the differentiated and non-differentiated subtypes. *Ann Epidemiol*, **13**, 24-31.
- Jemal A, Bray F, Center MM, et al (2011). Global cancer statistics. *CA Cancer J Clin*, **61**, 69-90.
- Jensen OM, Parkin DM, Lennan R, et al (1991). Cancer registration. Principles and Methods. Lyon (France): International Agency for Research on Cancer.
- Levi F, Lucchini F, Gonzalez JR, et al (2004). Monitoring falls in gastric cancer mortality in Europe. *Ann Oncol*, **15**, 338-45.
- Levi F, Lucchini F, Negri E, et al (2003). Mortality from major cancer sites in the European Union, 1955-1998. *Ann Oncol*, **14**, 490-5.
- Lee JK, Inoue M, Otani T, et al (2006). Gastric cancer screening and subsequent risk of gastric cancer: a large-scale population-based cohort study, with a 13-year follow-up in Japan. *Int J Cancer*, **118**, 2315-21.
- Look M, Gao F, Low CH, and Nambiar R (2001). Gastric cancer in Singapore. *Gastric Cancer*, **4**, 219-22.
- Maskarinec G, Noh JJ (2004). The effect of migration on cancer incidence among Japanese in Hawaii. *Ethn Dis*, **14**, 431-9.
- Munoz N, Correa P, Cuello C, Duque E (1968). Histologic types of gastric carcinoma in high- and low-risk areas. *Int J Cancer*, **3**, 809-18.
- Munoz N, Franceschi S (1997). Epidemiology of gastric cancer and perspectives for prevention. *Salud Publica Mex*, **39**, 318-30.
- Ngoan LT, Mizoue T, Fujino Y, et al 2002. Dietary factors and stomach cancer mortality. *Br J Cancer*, **87**, 37-42.
- Parkin DM (2004). International variation. *Oncogene*, **23**, 6329-40.
- Parkin DM, Whelan SL, Ferlay J, Teppo L, Thomas DB (eds) (2002). Cancer Incidence in Five Continents, Vol VIII, IARC Scientific Publications No. 155 IARC: Lyon, France.
- World Health Organization. International Classification of Diseases (ICD). Available at: <http://www.who.int/classifications/icd/en/>
- Yang L (2006). Incidence and mortality of gastric cancer in China. *World J Gastroenterol*, **12**, 17-20.