

Neonatal tetanus – report of a case

Milena Ilic¹, Ljiljana Pejic², Branislav Todorovic³, Bajram Hasani⁴, Sandra Stankovic², Dejan Milojevic², Daniela Djordjevic², Jelena Vucic², Zaklina Milosevic², Mitat Sahiti⁵, Goran Ristic⁶

¹Department of Epidemiology, University of Kragujevac Faculty of Medicine, and ²Clinic for Children's Internal Diseases, and ³Department of Epidemiology, University of Nish Faculty of Medicine, and Health Care Centers, ⁴Bujanovac and ⁵Presevo, and ⁶ Institute of Public Health, Vranje, Serbia

SUMMARY: Ilic M, Pejic L, Todorovic B, Hasani B, Stankovic S, Milojevic D, Djordjevic D, Vucic J, Milosevic Z, Sahiti M, Ristic G. Neonatal tetanus: report of a case. Turk J Pediatr 2010; 52: 404-408.

Neonatal tetanus is a severe, often fatal disease caused by the toxin *Clostridium tetani*. Neonatal tetanus is a generalized tetanus, which occurs in a neonate between 3-28 days of life. The findings indicated that tetanus in a newborn of an unvaccinated mother occurred after the application of non-sterile clay to the umbilical cord. This case was a seven-day-old male baby with progressive difficulty in feeding, trismus, hypertonicity, opisthotonos, and heart murmur. The patient was afebrile and eupneic, and had a history of non-sterile home delivery. In the past, the area of Bujanovac, Medvedja and Presevo had been exposed to mass immigration (especially due to the war in the territory of former Yugoslavia), which caused a serious problem for general practitioners, who had to be vigilant and ensure that all patients registered in their practice were fully immunized. This case has provided a clear indication of the necessity for strategies of both vaccination and ensuring hygienic conditions throughout pregnancy and delivery to prevent neonatal tetanus.


Key words: neonatal tetanus, infection, death, case report.

Neonatal tetanus is a severe, often fatal disease caused by the toxin *Clostridium tetani*, a ubiquitous spore-forming bacterium found in high concentrations in the soil and animal excrements (including human beings)^{1,2}. Neonatal tetanus is a generalized tetanus, which occurs in a neonate between 3-28 days of life, and is sometimes referred to as the disease of the seventh day^{2,3}. The newborn usually exhibits irritability, poor feeding, rigidity, facial grimacing, and severe spasms of touch²⁻⁴. Characteristic features are early spasms of the facial muscles (trismus or “lock-jaw” and “risus sardonicus”) followed by the spasm of the back muscles (opisthotonos) and sudden, generalized tonic seizures (tetanospasms)⁵⁻⁸. Glottis spasm, respiratory failure, and autonomic instability can result in death⁵.

During 2007, out of 17,012 tetanus cases reported worldwide, 6067 occurred in a neonate^{1,7}. Tetanus cases primarily occur in underdeveloped countries and account for up to

one-half of all neonatal deaths. Although easily prevented by maternal immunization with tetanus toxoid vaccine and aseptic obstetric and postnatal umbilical cord care practices, maternal and neonatal tetanus persists as a public health problem in 48 countries, mainly in Asia and Africa. Of the estimated 28 countries in the world that account for 90% of neonatal cases, 16 are in the African region^{1,8}. In these countries, activities are being undertaken to eliminate the disease in the near future.

Neonatal tetanus is associated with non-sterile delivery and umbilical cord care of newborns whose mothers do not have sufficient antitoxin levels to protect the neonates by transplacental transfer of maternal antibody^{6,8-10}. Risk factors for neonatal tetanus incidence are related to prenatal (lack of antenatal care for the pregnant women in a health facility, failure of immunization with tetanus toxoid), perinatal (delivery at home, births followed by untrained persons, failure of simple measures



crying right away and was cyanotic. The first suckle was 2-3 hours afterwards. He sucked the breast normally for 5 days, when suddenly he developed disinclination to sucking with braking of the jaws; he could not open his mouth and had difficulty opening the eyelids.

The mother, a 37-year-old woman born in Serbia, had no certificates of immunization, and it was therefore presumed that she had never been vaccinated. All thirteen births were at home. The first child died in the fourth month of life, the third child in the eighth and the fourth child in the tenth month. The causes of death of these children were unknown. The fifth child died presenting all symptoms and signs like the current patient. The other eight children, the patient's elder brothers and sisters, were healthy and had immunization following the normal schedule.

A low-income family, they lived at home without a supply network; they were supplied with drinking water from a public drink fountain. Both of the parents were unqualified and unemployed. They had a state health insurance.

Previously, the area of Bujanovac, Medvedja and Presevo had been exposed to mass immigration (especially due to the war in the territory of former Yugoslavia), which caused a serious problem for general practitioners, who had to be vigilant and ensure that all patients registered in their practice were fully immunized.

Discussion

In Serbia, neonatal tetanus is rare. The last two cases were reported in 1997¹⁷. This decline is associated with improvements in birth practices and increased level of population immunity following the initiation of routine tetanus toxoid vaccination since 1951.

Neonatal tetanus results from cord contamination during unsanitary delivery conditions, coupled with a lack of maternal immunization.

In newborns, the common nidus of infection is the umbilical cord, especially a septic umbilicus or any superficial wound; in many cases, it may not be detectable^{15,18}. Most cases follow an acute injury, such as a puncture

wound, a laceration or an abrasion. Although the use of a new razor was found to be significantly protective for neonatal tetanus appearance^{10,11}, this association has been refuted in some studies^{19,20}. Home delivery, the mother's education, a cleaned cutting tool, the application of antibiotics at delivery, and hand washing by the delivery attendant remained protective¹¹⁻¹⁴.

In this case, the mother cut the umbilical cord with a new, but non-sterile razor and tied it with knitting string. The umbilical cord was very poorly treated and the umbilicus showed signs of inflammation on admission, when omphalitis was noticed. In this newborn, the nidus of infection was probably a septic umbilicus or any superficial wound (microabrasion) around the cord, as was noticed in other reports^{11,20,21}. The spores need tissue with the proper anaerobic conditions to germinate; the ideal media are wounds with tissue necrosis. *Clostridium tetani* is recovered from wounds in only some 30% of cases, and the organism is sometimes isolated from patients who do not have tetanus^{5,22}. In this case, *Clostridium tetani* was not revealed, but culture from the umbilical cord grew several aerobic bacterial species (*Staphylococcus aureus* and *Streptococcus beta haemolyticus* B). This active infection, such as those with dead or devitalized tissue, was ideal for germination of the spores and release of toxin. Intermediate determinants of omphalitis may have included hygiene-related practices.

Data analysis suggests that the main source of *Clostridium tetani* may be the hands of the birth attendant, while the main mode of contamination may be the dressing of the wound stump¹⁰⁻¹⁴. In this case, poor personal hygiene maintenance, including type of birthing surface, cord care (tying, cutting, topical applications), infant bathing practices, attendants' hand washing practices, and skin-to-skin contact between mother and newborn were noticed.

Mothers with previous history of neonatal tetanus babies were shown to have a significantly increased risk, and accounted for more than one-third of all cases^{19,23,24}. In this case, one child died presenting all the symptoms and signs as observed in the current patient. This may indicate the importance of both poor hygienic condition and lack of mother's immunization.

REFERENCES

1. World Health Organization. Vaccines and Biologicals. WHO-recommended standards for surveillance of selected vaccine-preventable diseases. Geneva: World Health Organization; 2003. Available at: <http://www.who.int/vaccines-documents/>
2. Centers for Disease Control and Prevention. Case definitions for infectious conditions under public health surveillance. *MMWR Recomm Rep* 1997; 46(No. RR-10): 1-55.
3. World Health Organization. Maternal and Neonatal Tetanus Elimination by 2005. Geneva: World Health Organization; 2005. Available at: <http://www.who.int/vaccines-documents/>
4. Centers for Disease Control and Prevention. Prevention of Pertussis, Tetanus, and Diphtheria Among Pregnant and Postpartum Women and Their Infants. Recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR Recomm Rep* 2008; 57: 1-47.
5. Cook TM, Protheroe RT, Handel JM. Tetanus: a review of the literature. *Br J Anaesth* 2001; 87: 477-487.
6. Roper MH, Vandelaer JH, Gasse FL. Maternal and neonatal tetanus. *Lancet* 2007; 370: 1947-1959.
7. World Health Organization. Global burden of disease and injury series. Volume IV. In: Murray CJ, Lopez AD, Mathers CD (eds). *The Global Epidemiology of Infectious Diseases*. Geneva: World Health Organization; 2004.
8. World Health Organization. Tetanus vaccine. *Wkly Epidemiol Rec* 2006; 81: 198-208.
9. Stanfield JP, Galazka A. Neonatal tetanus in the world today. *Bull World Health Organ* 1984; 62: 647-669.
10. Bennett J, Schooley M, Traverso H, Bano Agha S, Boring J. Bundling, a newly identified risk factor for neonatal tetanus: implications for global control. *Int J Epidemiol* 1996; 25: 879-884.
11. Parashar UD, Bennett JV, Boringa JR, Hladyd WG. Topical antimicrobials applied to the umbilical cord stump: a new intervention against neonatal tetanus. *Int J Epidemiol* 1998; 27: 904-908.
12. Hlady WG, Bennett JV, Samadi AR, et al. Neonatal tetanus in rural Bangladesh: risk factors and toxoid efficacy. *Am J Public Health* 1992; 82: 1365-1369.
13. Leroy O, Garenne M. Risk factors of neonatal tetanus in Senegal. *Int J Epidemiol* 1991; 20: 521-526.
14. Raza SA, Akhtar S, Avan BI, Hamza H, Rahbar MH. A matched case-control study of risk factors for neonatal tetanus in Karachi, Pakistan. *J Postgrad Med* 2004; 50: 247-251.
15. Davies-Adetugbo AA, Torimiro SE, Ako-Nai KA. Prognostic factors in neonatal tetanus. *Trop Med Int Health* 1998; 3: 9-13.
16. Quddus A, Luby S, Rahbar M, Pervaiz Y. Neonatal tetanus: mortality rate and risk factors in Loralai District, Pakistan. *Int J Epidemiol* 2002; 31: 648-653.

17. World Health Organization. WHO vaccine-preventable diseases: monitoring system 2009 global summary. Available from URL: http://www.who.int/immunization_monitoring/
18. Centers for Disease Control and Prevention. Tetanus-United States, 1985-6. *MMWR Morb Mortal Wkly Rep* 1987; 36: 477-481.
19. Traverso HP, Kamil S, Rahim H, Samadi AR, Boring JR, Bennett JV. A reassessment of risk factors for neonatal tetanus. *Bull World Health Organ* 1991; 69: 573-579.
20. Mullany LC, Darmstadt GL, Katz J, et al. Risk factors for umbilical cord infection among newborns of southern Nepal. *Am J Epidemiol* 2007; 165: 203-211.
21. Bjerregaard P, Steinglass R, Mutie DM, Kimani G, Mjomba M, Orinda V. Neonatal tetanus mortality in coastal Kenya: a community survey. *Int J Epidemiol* 1993; 22: 163-169.
22. Guardiola A, Teixeira AM, da Silva CA, et al. Neonatal tetanus. *J Pediatr (Rio J)* 2000; 76: 391-394.
23. Chai F, Prevots DR, Wang X, Birmingham M, Zhang R. Neonatal tetanus incidence in China, 1996-2001, and risk factors for neonatal tetanus, Guangxi Province, China. *Int J Epidemiol* 2004; 33: 551-557.
24. Bennett J, Macia J, Traverso H, Banoagha S, Malooly C, Boring J. Protective effects of topical antimicrobials against neonatal tetanus. *Int J Epidemiol* 1997; 26: 897-903.
25. Deming MS, Rongou JB, Kristiansen M, et al. Tetanus toxoid coverage as an indicator of serological protection against neonatal tetanus. *Bull World Health Organ* 2002; 80: 696-703.
26. Dietz V, Galazka A, van Loon F, Cochi S. Factors affecting the immunogenicity and potency of tetanus toxoid: implications for the elimination of neonatal and non-neonatal tetanus as public health problems. *Bull World Health Organ* 1997; 75: 81-93.
27. Dietz V, Milstien JB, van Loon F, Cochi S, Bennett J. Performance and potency of tetanus toxoid: implications for eliminating neonatal tetanus. *Bull World Health Organ* 1996; 74: 619-628.
28. World Health Organization. Validation of neonatal tetanus elimination in Andhra Pradesh, India. *Wkly Epidemiol Rec* 2004; 79: 292-297.
29. World Health Organization. Field Guide for Surveillance of Vaccine Preventable Diseases. Nepal: World Health Organization; 2005. Available at: <http://www.whoipd.org/>