

ORIGINAL ARTICLE

Mortality from cancer of the lung in Serbia

M. Ilic¹, H. Vlajinac², J. Marinkovic³, Z. Blazic⁴

¹Department of Epidemiology, Faculty of Medical Sciences, University of Kragujevac, Kragujevac; ²Institute of Epidemiology, Faculty of Medicine, University of Belgrade, Belgrade; ³Institute of Medical Statistics and Informatics, Faculty of Medicine, University of Belgrade, Belgrade; ⁴Clinic of Otorhinolaryngology and Maxillofacial Surgery, Clinical Center of Serbia, Belgrade, Serbia

Summary

Purpose: To estimate the death rates for lung cancer and their secular trends in the population of Serbia, excluding the autonomous province of Kosovo and Metohia, over the 1991-2009 period.

Methods: A descriptive epidemiological method was employed. Trend of the lung cancer mortality rates was estimated using joinpoint linear regression analysis. An average annual percentage of change (AAPC) was computed for trend using linear models assuming a Poisson distribution, and the corresponding 95% confidence interval (CI).

Results: The mortality rate from lung cancer in Serbia ranks as the highest in the world, and it has been increasing continuously from 1991 (AAPC = + 1.9; 95% CI=1.7-2.2). A significant increase in mortality was present in both the male population (AAPC = + 1.4; 95% CI=1.2-1.6), and the female population (AAPC = + 3.9; 95% CI=3.6-4.3). How-

ever, a significant decline in lung cancer mortality in men was seen in some age groups. In young men (35-39 and 40-44 years age groups), lung cancer death rates decreased continuously from 1991, by - 5.1% and - 2.6% per year, respectively. Among men in the 45-49 years age group, a marked increase of lung cancer mortality was observed from 1991 to 1998 (by + 6.5% per year), followed by significant decrease (- 1.9% per year). Among women, only in the youngest age group (35-39 years) a declining trend was present (- 0.6% per year), yet without significance.

Conclusion: Lung cancer mortality rates in Serbia indicate the importance of consistent application of measures of primary and secondary prevention that have been proven effective in other countries.

Key words: descriptive epidemiological study, joinpoint analysis, lung cancer, mortality

Introduction

Lung cancer was the leading cause of cancer-related deaths in males in 2008 globally, and the second leading cause of cancer-related deaths among females [1]. Lung cancer ranks nearly fifth of all cancer deaths worldwide, with significant geographic variations in frequency and distribution [1-3].

The mortality rate from lung cancer in Serbia for males and females taken together ranks third in the world, after the rates in Hungary and French Polynesia [2]. Lung cancer mortality is more than twice as frequent in men than in women [3,4]. In males, the highest lung cancer death rates are in Central and Eastern Europe, and then North America, Eastern Asia and Western Europe

[2]. In females, the highest lung cancer death rates are found in North America, North Europe, and Eastern Asia.

Since the 1980s, lung cancer mortality rates have been declining in most developed countries, but not in all of them [4-6]. Despite the persistent decline of lung cancer in males in the United States of America and the European Union (each by - 1.9% per year), the trends of lung cancer mortality in females continuously rise (by + 0.3% and + 1.7% per year, respectively) [5,7]. It is considered that the major contribution for the decrease in lung cancer mortality rate is attributable to reduction in tobacco smoking [8,9].

The aim of the present study was to estimate death rates for lung cancer in the population of Serbia and their trends during the period 1991-2009.




Table 2. Joinpoint regression analysis of lung cancer mortality in Serbia by gender and age in 1991-2009, excluding the autonomous province of Kosovo and Metohia

Age [§]	1991-2009		Trend 1		Trend 2	
	Average age-specific rates (per 100,000)	AAPC (95%CI)	Years	APC (95%CI)	Years	APC (95%CI)
Males						
35-39	7.4	- 5.1* (-7.4 to -2.8)				
40-44	24.8	- 2.6* (-3.8 to -1.4)				
45-49	62.7	+ 1.3* (0.4-2.2)	1991-1998	+ 6.5* (4.5-8.5)	1998-2009	-1.9* (-2.8 to -0.9)
50-54	120.1	+ 1.1 (-0.8-3.0)	1991-2007	+ 2.3* (1.7-3.0)	2007-2009	- 8.4 (-23.1-9.1)
55-59	184.6	+ 2.2* (1.2-3.2)	1991-1998	- 0.4 (-2.5-1.7)	1998-2009	+ 3.9* (2.8-5.0)
60-64	260.7	+ 1.0* (0.1-1.9)	1991-2001	- 0.8 (-1.9-0.4)	2001-2009	+ 3.2* (1.5-5.0)
65-69	324.5	+ 0.8* (0.5-1.2)				
70-74	352.8	+ 2.0* (1.6-2.4)				
75-79	322.3	+ 3.0* (2.2-3.8)				
80-84	234.6	+ 3.7* (2.6-4.8)				
85+	137.8	+ 3.0* (0.7-5.3)				
All males		+ 1.4* (1.2-1.6)				
Females						
35-39	3.3	- 0.6 (-4.6-3.7)				
40-44	9.6	+ 1.8 (-0.1-3.8)				
45-49	17.4	+ 4.3* (2.7-5.9)				
50-54	31.0	+ 6.4* (5.4-7.3)				
55-59	40.3	+ 5.2* (3.7-6.9)				
60-64	48.7	+ 4.1* (3.3-5.0)				
65-69	45.2	+ 3.0* (2.1-3.8)				
70-74	71.8	+ 3.6* (2.7-4.4)				
75-79	74.5	+ 3.3* (2.4-4.1)				
80-84	65.0	+ 3.6* (2.1-5.2)				
85+	46.4	+ 1.9 (-0.3-4.2)				
All females		+ 3.9* (3.6-4.3)				


* Joinpoint significantly different from zero at $\alpha = 0.05$; [§] Joinpoint results are not shown for the age subgroup ≤ 34 , because there were less than 10 cases in any year.

AAPC: average annual percent change, APC: annual percent change, CI: confidence interval

decreased continuously from 1991, by - 5.1% and - 2.6% per year, respectively. In men 65 years and older, age specific mortality rates increased continuously over the entire study period. In middle-aged men (45-64 ages) two patterns of changes in lung cancer mortality were observed. Among men in the 45-49 age group, a marked increase of lung cancer mortality from 1991 to 1998 (by + 6.5% per year) was followed by significant decrease (- 1.9% per year). In men 50-54 years old, after a significant increase from 1991 to 2007 (+ 2.3 % per year), lung cancer death rates began to decline in the last 3 years of the period ob-

served, but not significantly (- 8.4 % per year). In men 55-59 and 60-64 years old nonsignificant decline of mortality rates (by - 0.4% and - 0.8% per year, respectively) were followed by significant increase from 1998 (+ 3.8% per year), and 2001, respectively (+ 3.2% per year).

Among women, in all age groups from 45 to 84 years, a significant rise in lung cancer death rates was observed continuously from 1991. Only in the youngest age group (35-39 years) a declining trend was present (- 0.6% per year), but without significance.



as well as in the European Union as a whole [15].

About 90% of lung cancers in men and 83% in women in developed countries are caused by smoking [16]. In Serbia, 82.8% of the lung cancer burden for men and 90.2% for women was attributable to smoking [17]. Levels and changes in the prevalence of smoking may explain both differences in lung cancer morbidity and mortality among countries and between men and women, and decreasing and increasing trend of lung cancer rates. The tobacco exposure in Serbia is still higher than in developed countries. According to data for the Belgrade population there were 49% male and 25% female smokers in the years 1976-1977, and 51% male and 37% female smokers in the years 1988-1989 [18]. Thanks to antismoking campaign which was intensified from 2000, in 2006 in Serbia 33.6% of the population were smokers (38.1 males vs 29.9% females), suggesting a reduction of the smoking rate by 6.9% in comparison with 2000, when the habit was 47.9% among men and

38.1% among women [19]. The highest percentage of smokers (46.9%) was recorded in the 35 to 44 years age group. In comparison to 2000, the number of young people aged 15-19 who have never tried to smoke was increased by 7.3%. Serbia joined World Health Organization convention on tobacco control and this convention became effective in June 2006. Taking into account the period of latency, one could expect that lung cancer mortality rates, especially in Serbian women, will continue to increase in the following years. Providing successful tobacco control strategy, the decreasing trend could be expected to begin in about 10 years. The already seen decline in lung cancer mortality rates in young men are promising.

Acknowledgements

This work was supported by the Ministry of Education and Science of Republic of Serbia, through Contract No. 175042.

References

1. Ferlay J, Shin HR, Bray F, Forman D, Mathers C, Parkin DM. Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *Int J Cancer* 2010; 127: 2893-2917.
2. IARC. World Cancer Report, 2008. Boyle P, Levin BE (Eds): IARC Press, Lyon. Available at: <http://monographs.iarc.fr/>. Accessed: August 16, 2012.
3. Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D. Global cancer statistics. *CA Cancer J Clin* 2011; 61: 69-90.
4. Jemal A, Thun MJ, Ries LA et al. Annual report to the nation on the status of cancer, 1975-2005, featuring trends in lung cancer, tobacco use, and tobacco control. *J Natl Cancer Inst* 2008; 100: 1672-1694.
5. Levi F, Lucchini F, Negri E, La Vecchia C. Continuing declines in cancer mortality in the European Union. *Ann Oncol* 2007; 18: 593-595.
6. Bray FI, Weiderpass E. Lung cancer mortality trends in 36 European countries: secular trends and birth cohort patterns by sex and region 1970-2007. *Int J Cancer* 2010; 126: 1454-1466.
7. Ries L, Harkins D, Krapcho M et al. SEER Cancer Statistics Review, 1975-2003. Bethesda, Maryland: National Cancer Institute, 2006. Available at: http://seer.cancer.gov/csr/1975_2003/results_merged/sect_15_lung_bronchus.pdf. Accessed: August 16, 2012.
8. Lopez AD, Collishaw NE, Piha T. A descriptive model of the cigarette epidemic in developed countries. *Tob Control* 1994; 3: 242-247.
9. Thun MJ, Burns DM. Health impact of "reduced yield" cigarettes: a critical assessment of the epidemiological evidence. *Tob Control* 2001; 10 (Suppl 1): i4-11.
10. Segi M, Fujisaku S, Kurihara M, Narai Y, Sasajima K. The age-adjusted death rates for malignant neoplasms in some selected sites in 23 countries in 1954-1955 and their geographical correlation. *Tohoku J Exp Med* 1960; 72: 91-103.
11. La Vecchia C, Bosetti C, Lucchini F et al. Cancer mortality in Europe, 2000-2004, and an overview of trends since 1975. *Ann Oncol* 2010; 21: 1323-1360.
12. Bosetti C, Bertuccio P, Levi F, Lucchini F, Negri E, La Vecchia C. Cancer mortality in the European Union, 1970-2003, with a joinpoint analysis. *Ann Oncol* 2008; 19: 631-640.
13. Shkolnikov VM, McKee M, Vallin J et al. Cancer mortality in Russia and Ukraine: validity, competing risks, and cohort effects. *Int J Epidemiol* 1999; 28: 19-29.
14. Cayuela A, Rodríguez-Domínguez S, López-Campos JL, Vigil E, Otero R. Lung cancer mortality trends in Spain between 1980 and 2005. *Arch Broncopneumol* 2008; 44: 70-74.
15. Bosetti C, Levi F, Lucchini F, Negri E, La Vecchia C. Lung cancer mortality in European women: recent trends and perspectives. *Ann Oncol* 2005; 16: 1597-1604.
16. Peto R, Lopez AD, Boreham J, Thun M. Mortality from smoking in developed countries 1950-2000 (2nd Edn; revised June 2006). Available from: www.deaths-fromsmoking.net. Accessed: August 16, 2012.
17. Atanasković-Marković Z, Bjegović V et al. Burden of disease and injury in Serbia. Belgrade: Republic of Serbia Ministry of Health; 2003. Available at: www.sbds.sr.gov.yu. Accessed: August 16, 2012.
18. Vlainjac H, Adanja B, Jarebinski M. Smoking habits of Belgrade urban population. *Med Invest* 1990; 23: 73-76.
19. Ministry of Health, Republic of Serbia. National Health Survey, Serbia 2006. Ministry of Health, Republic of Serbia, 2007, Belgrade.