

**In vitro evaluation of the genotoxicity of ritodrine and verapamil in human lymphocytes**

O. Milosevic-Djordjevic, D. Grujicic, G. Joksic and D. Marinkovic

*Hum Exp Toxicol* 2011 30: 398 originally published online 20 May 2010

DOI: 10.1177/0960327110372404

The online version of this article can be found at:

<http://het.sagepub.com/content/30/5/398>

---

Published by:



<http://www.sagepublications.com>

**Additional services and information for *Human & Experimental Toxicology* can be found at:**

**Email Alerts:** <http://het.sagepub.com/cgi/alerts>

**Subscriptions:** <http://het.sagepub.com/subscriptions>

**Reprints:** <http://www.sagepub.com/journalsReprints.nav>

**Permissions:** <http://www.sagepub.com/journalsPermissions.nav>

**Citations:** <http://het.sagepub.com/content/30/5/398.refs.html>

# **In vitro evaluation of the genotoxicity of ritodrine and verapamil in human lymphocytes**

**O Milošević-Djordjević<sup>1,2</sup>, D Grujičić<sup>1</sup>, G Joksić<sup>3</sup> and D Marinković<sup>4</sup>**

## **Abstract**

The aim of this study was to investigate the genotoxic effects of ritodrine and verapamil on human peripheral lymphocytes in vitro using micronucleus (MN) test. Also, fluorescence in situ hybridization (FISH) with a centromeric probe was performed to determine the origin of the induced MN. Cells were treated with  $8.4 \times 10^{-6}$  M –  $25.2 \times 10^{-4}$  M concentrations for ritodrine and  $0.56$ – $1.1 \times 10^{-5}$  M concentrations for verapamil, separately and combined. The MN frequencies showed increase after all treatments, but the difference between treated cells and untreated controls were found to be statistically significant only in the concentration range from  $8.4 \times 10^{-5}$  M –  $4.5 \times 10^{-4}$  M for ritodrine,  $1.1$ – $3.3 \times 10^{-5}$  M for verapamil, and in combined treatment with concentrations  $8.4 \times 10^{-5}$  M +  $1.1 \times 10^{-5}$  M for ritodrine and verapamil. The highest tested concentrations of both medicaments showed cytotoxic effect. Both medicaments decreased the nuclear division index (NDI) in tested concentrations. The results of FISH analysis suggest that verapamil, separately or combined with ritodrine, shows to a larger extent aneugenic than clastogenic effect.

## **Keywords**

micronucleus, lymphocytes, in vitro, ritodrine, verapamil

## **Introduction**

Micronuclei (MN) are small, round chromatin bodies in the cytoplasm of cells that originate from cell division as the result of the elimination of acentric chromosome fragments or whole chromosomes. There are several genotoxic studies on pharmaceutical compounds using this assay.<sup>1–3</sup> To distinguish between MN derived from acentric fragments and MN containing whole chromosomes, fluorescence in situ hybridization (FISH) with centromere-specific DNA probes is being increasingly used in monitoring studies,<sup>3,4</sup> and the presence of hybridization signal in an MN is a direct measure of the presence of a centromere.<sup>5</sup>

Ritodrine hydrochloride (ritodrine), IUPAC name 4-[2-[2-hydroxy-2-(4-hydroxyphenyl)-1-methyl-ethyl] aminoethyl] phenol, belongs to  $\beta_2$  group of selective agonists that have special affinity for  $\beta_2$  receptors in myometrium (Figure 1). Stimulating these receptors, they induce uterus relaxation, especially during gravidity. This property is of prominent importance for the prevention and protection from premature

contractions of the uterus in gravidity.<sup>6</sup> Ritodrine can stop the threatening abortion or postpone premature labor. It is administered orally or intravenously. Oral dose of  $3 \times 40$  mg in combination with intravenous transfusion of  $2 \times 50$  mg per day is administered until the uterus contractions stop, to be followed by oral administration only. In addition to the heart stimulation (tachycardia, more forceful heart contractions and irritability), ritodrine also induces hypotension, tremor at rest, anxiety, dizziness and nausea.<sup>7</sup>

<sup>1</sup> Faculty of Science, University of Kragujevac, Kragujevac, Serbia

<sup>2</sup> Medical Faculty, University of Kragujevac, Kragujevac, Serbia

<sup>3</sup> "Vinča" Institute of Nuclear Sciences, Belgrade, Serbia

<sup>4</sup> Serbian Academy of Sciences and Arts, Belgrade, Serbia

## **Corresponding author:**

Olivera Milošević-Djordjević, Faculty of Science, University of Kragujevac, Institute of Biology and Ecology, Department of Genetics, Radoja Domanovića 12, P.O. Box 60 34000, Kragujevac, Serbia  
Email: olivera@kg.ac.rs















